



On the Analytic Part of Univalent Harmonic Mappings

Iigiz R. Kayumov¹ · Saminthan Ponnusamy² ·
Le Anh Xuan¹

Received: 3 February 2017 / Accepted: 18 December 2017 / Published online: 22 January 2018
© Springer International Publishing AG, part of Springer Nature 2018

Abstract In this article we obtain two sharp results concerning the analytic part of harmonic mappings $f = h + \bar{g}$ from the class $S_H^0(\mathcal{S})$ which was recently introduced by Ponnusamy and Sairam Kaliraj. For example, we get the sharp estimate for $|\arg h'(z)|$ in the case when $|z| \leq 1/\sqrt{2}$ and obtain the sharp radius of convexity for h . Our approach is applicable to a more general situation. Finally, we determine simple condition on the analytic part of univalent harmonic mappings so that it is in H_p spaces for $0 < p < 1/3$.

Keywords Harmonic univalent and convex mappings · Rotation theorem · Schwarz–Pick Lemma · Koebe transform · Disk automorphism · Harmonic (analytic) Hardy spaces · Integral means

Mathematics Subject Classification Primary 30C45 · 30C50 · 30C55 · 30H10 · 31A05; Secondary 30C75 · 31A20

Communicated by David Shoikhet.

✉ Saminthan Ponnusamy
samy@isichennai.res.in; samy@iitm.ac.in

Iigiz R. Kayumov
ikayumov@kpfu.ru

Le Anh Xuan
laxuan@ctu.edu.vn

¹ Kazan Federal University, Kremlevskaya 18, Kazan, Russia 420 008

² Department of Mathematics, Indian Institute of Technology Madras, Chennai 600 036, India